Rend. Istit. Mat. Univ. Trieste
Volume 44 (2012), 413–422

Rank two globally generated vector bundles with $c_1 \leq 5$

Ludovica Chiodera and Philippe Ellia

Abstract. *We classify globally generated rank two vector bundles on $\mathbb{P}^n$, $n \geq 3$, with $c_1 \leq 5$. The classification is complete but for one case ($n = 3$, $c_1 = 5$, $c_2 = 12$).*

Keywords: Vector bundles, rank two, globally generated, projective space
MS Classification 2010: 14F05, 14M15

1. Introduction.

Vector bundles generated by global sections are basic objects in projective algebraic geometry. Globally generated line bundles correspond to morphisms to a projective space, more generally higher rank bundles correspond to morphism to (higher) Grassmann varieties. For this last point of view (that won't be touched in this paper) see [10, 12, 13]. Also globally generated vector bundles appear in a variety of problems ([7] just to make a single, recent example).

In this paper we classify globally generated rank two vector bundles on $\mathbb{P}^n$ (projective space over k, $\overline{k} = k$, $ch(k) = 0$), $n \geq 3$, with $c_1 \leq 5$. The result is:

Theorem 1.1. *Let E be a rank two vector bundle on $\mathbb{P}^n$, $n \geq 3$, generated by global sections with Chern classes c_1, c_2, $c_1 \leq 5$.*

1. *If $n \geq 4$, then E is the direct sum of two line bundles*
2. *If $n = 3$ and E is indecomposable, then*

$$(c_1, c_2) \in S = \{((2,2),(4,5),(4,6),(4,7),(4,8),(5,8),(5,10),(5,12)\}.$$

If E exists there is an exact sequence:

$$0 \to \mathcal{O} \to E \to \mathcal{I}_C(c_1) \to 0 \ (*)$$

where $C \subset \mathbb{P}^3$ is a smooth curve of degree c_2 with $\omega_C(4 - c_1) \simeq \mathcal{O}_C$. The curve C is irreducible, *except maybe if $(c_1, c_2) = (4, 8)$: in this case C can be either irreducible or the disjoint union of two smooth conics.*

3. *For every* $(c_1, c_2) \in S$, $(c_1, c_2) \neq (5, 12)$, *there exists a rank two vector bundle on* $\mathbb{P}^3$ *with Chern classes* (c_1, c_2) *which is globally generated (and with an exact sequence as in 2.).*

The classification is complete, but for one case: we are unable to say if there exist or not globally generated rank two vector bundles with Chern classes $c_1 = 5, c_2 = 12$ on $\mathbb{P}^3$.

2. Rank two vector bundles on $\mathbb{P}^3$.

2.1. General facts.

For completeness let's recall the following well known results:

LEMMA 2.1. *Let* E *be a rank* r *vector bundle on* $\mathbb{P}^n$, $n \geq 3$. *Assume* E *is generated by global sections.*

1. *If* $c_1(E) = 0$, *then* $E \simeq r.\mathcal{O}$

2. *If* $c_1(E) = 1$, *then* $E \simeq \mathcal{O}(1) \oplus (r-1).\mathcal{O}$ *or* $E \simeq T(-1) \oplus (r-n).\mathcal{O}$.

Proof. If $L \subset \mathbb{P}^n$ is a line then $E|L \simeq \bigoplus_{i=1}^r \mathcal{O}_L(a_i)$ by a well known theorem and $a_i \geq 0, \forall i$ since E is globally generated. It turns out that in both cases: $E|L \simeq \mathcal{O}_L(c_1) \oplus (r-1).\mathcal{O}_L$ for every line L, i.e. E is uniform. Then 1. follows from a result of Van de Ven ([14]), while 2. follows from IV. Prop. 2.2 of [4]. □

LEMMA 2.2. *Let* E *be a rank two vector bundle on* $\mathbb{P}^n$, $n \geq 3$. *If* E *has a nowhere vanishing section then* E *splits. If* E *is generated by global sections and doesn't split then* $h^0(E) \geq 3$ *and a general section of* E *vanishes along a smooth curve,* C, *of degree* $c_2(E)$ *such that* $\omega_C(4 - c_1) \simeq \mathcal{O}_C$. *Moreover* $\mathcal{I}_C(c_1)$ *is generated by global sections.*

LEMMA 2.3. *Let* E *be a non split rank two vector bundle on* $\mathbb{P}^3$ *with* $c_1 = 2$. *If* E *is generated by global sections then* E *is a null-correlation bundle.*

Proof. We have an exact sequence: $0 \to \mathcal{O} \to E \to \mathcal{I}_C(2) \to 0$, where C is a smooth curve with $\omega_C(2) \simeq \mathcal{O}_C$. It follows that C is a disjoint union of lines. Since $h^0(\mathcal{I}_C(2)) \geq 2$, $d(C) \leq 2$. Finally $d(C) = 2$ because E doesn't split. □

This settles the classification of rank two globally generated vector bundles with $c_1(E) \leq 2$ on $\mathbb{P}^3$.

2.2. Globally generated rank two vector bundles with $c_1 = 3$.

The following result has been proved in [10] (with a different and longer proof).

PROPOSITION 2.4. *Let E be a rank two globally generated vector bundle on $\mathbb{P}^3$. If $c_1(E) = 3$ then E splits.*

Proof. Assume a general section vanishes in codimension two, then it vanishes along a smooth curve C such that $\omega_C \simeq \mathcal{O}_C(-1)$. Moreover $\mathcal{I}_C(3)$ is generated by global sections. We have $C = \cup_{i=1}^r C_i$ (disjoint union) where each C_i is smooth irreducible with $\omega_{C_i} \simeq \mathcal{O}_{C_i}(-1)$. It follows that each C_i is a smooth conic. If $r \geq 2$ let $L = \langle C_1 \rangle \cap \langle C_2 \rangle$ ($\langle C_i \rangle$ is the plane spanned by C_i). Every cubic containing C contains L (because it contains the four points $C_1 \cap L$, $C_2 \cap L$). This contradicts the fact that $\mathcal{I}_C(3)$ is globally generated. Hence $r = 1$ and $E = \mathcal{O}(1) \oplus \mathcal{O}(2)$. □

2.3. Globally generated rank two vector bundles with $c_1 = 4$.

Let's start with a general result:

LEMMA 2.5. *Let E be a non split rank two vector bundle on $\mathbb{P}^3$ with Chern classes c_1, c_2. If E is globally generated and if $c_1 \geq 4$ then:*

$$c_2 \leq \frac{2c_1^3 - 4c_1^2 + 2}{3c_1 - 4}.$$

Proof. By our assumptions a general section of E vanishes along a smooth curve, C, such that $\mathcal{I}_C(c_1)$ is generated by global sections. Let U be the complete intersections of two general surfaces containing C. Then U links C to a smooth curve, Y. We have $Y \neq \varnothing$ since E doesn't split. The exact sequence of liaison: $0 \to \mathcal{I}_U(c_1) \to \mathcal{I}_C(c_1) \to \omega_Y(4 - c_1) \to 0$ shows that $\omega_Y(4 - c_1)$ is generated by global sections. Hence $\deg(\omega_Y(4 - c_1)) \geq 0$. We have $\deg(\omega_Y(4 - c_1)) = 2g' - 2 + d'(4 - c_1)$ ($g' = p_a(Y)$, $d' = \deg(Y)$). So $g' \geq \frac{d'(c_1-4)+2}{2} \geq 0$ (because $c_1 \geq 4$). On the other hand, always by liaison, we have: $g' - g = \frac{1}{2}(d' - d)(2c_1 - 4)$ ($g = p_a(C)$, $d = \deg(C)$). Since $d' = c_1^2 - d$ and $g = \frac{d(c_1-4)}{2} + 1$ (because $\omega_C(4 - c_1) \simeq \mathcal{O}_C$), we get: $g' = 1 + \frac{d(c_1-4)}{2} + \frac{1}{2}(c_1^2 - 2d)(2c_1 - 4) \geq 0$ and the result follows. □

Now we have:

PROPOSITION 2.6. *Let E be a rank two globally generated vector bundle on $\mathbb{P}^3$. If $c_1(E) = 4$ and if E doesn't split, then $5 \leq c_2 \leq 8$ and there is an exact sequence: $0 \to \mathcal{O} \to E \to \mathcal{I}_C(4) \to 0$, where C is a smooth irreducible elliptic curve of degree c_2 or, if $c_2 = 8$, C is the disjoint union of two smooth elliptic quartic curves.*

Proof. A general section of E vanishes along C where C is a smooth curve with $\omega_C = \mathcal{O}_C$ and where $\mathcal{I}_C(4)$ is generated by global sections. Let $C = C_1 \cup ... \cup C_r$ be the decomposition into irreducible components: the union is disjoint, each C_i is a smooth elliptic curve hence has degree at least three.
By Lemma 2.5 $d = \deg(C) \leq 8$. If $d \leq 4$ then C is irreducible and is a complete intersection which is impossible since E doesn't split. If $d = 5$, C is smooth irreducible.
Claim: If $8 \geq d \geq 6$, C cannot contain a plane cubic curve.
Assume $C = P \cup X$ where P is a plane cubic and where X is a smooth elliptic curve of degree $d - 3$. If $d = 6$, X is also a plane cubic and every quartic containing C contains the line $\langle P \rangle \cap \langle X \rangle$. If $deg(X) \geq 4$ then every quartic, F, containing C contains the plane $\langle P \rangle$. Indeed $F|H$ vanishes on P and on the $deg(X) \geq 4$ points of $X \cap \langle P \rangle$, but these points are not on a line so $F|H = 0$. In both cases we get a contradiction with the fact that $\mathcal{I}_C(4)$ is generated by global sections. The claim is proved.
It follows that, if $8 \geq d \geq 6$, then C is irreducible except if $C = X \cup Y$ is the disjoint union of two elliptic quartic curves. □

Now let's show that all possibilities of Proposition 2.6 do actually occur. For this we have to show the existence of a smooth irreducible elliptic curve of degree d, $5 \leq d \leq 8$ with $\mathcal{I}_C(4)$ generated by global sections (and also that the disjoint union of two elliptic quartic curves is cut off by quartics).

LEMMA 2.7. *There exist rank two vector bundles with $c_1 = 4, c_2 = 5$ which are globally generated. More precisely any such bundle is of the form $\mathcal{N}(2)$, where $\mathcal{N}$ is a null-correlation bundle (a stable bundle with $c_1 = 0, c_2 = 1$).*

Proof. The existence is clear (if $\mathcal{N}$ is a null-correlation bundle then it is well known that $\mathcal{N}(k)$ is globally generated if $k \geq 1$). Conversely if E has $c_1 = 4, c_2 = 5$ and is globally generated, then E has a section vanishing along a smooth, irreducible quintic elliptic curve (cf 2.6). Since $h^0(\mathcal{I}_C(2)) = 0$, E is stable, hence $E = \mathcal{N}(2)$. □

LEMMA 2.8. *There exist smooth, irreducible elliptic curves, C, of degree 6 with $\mathcal{I}_C(4)$ generated by global sections.*

Proof. Let X be the union of three skew lines. The curve X lies on a smooth quadric surface, Q, and has $\mathcal{I}_X(3)$ globally generated (indeed the exact sequence $0 \to \mathcal{I}_Q \to \mathcal{I}_X \to \mathcal{I}_{X,Q} \to 0$ twisted by $\mathcal{O}(3)$ reads like: $0 \to \mathcal{O}(1) \to \mathcal{I}_C(3) \to \mathcal{O}_Q(3,0) \to 0$). The complete intersection, U, of two general cubics containing X links X to a smooth curve, C, of degree 6 and arithmetic genus 1. Since, by liaison, $h^1(\mathcal{I}_C) = h^1(\mathcal{I}_X(-2)) = 0$, C is irreducible. The exact sequence of liaison: $0 \to \mathcal{I}_U(4) \to \mathcal{I}_C(4) \to \omega_X(2) \to 0$ shows that $\mathcal{I}_C(4)$ is globally generated. □

In order to prove the existence of smooth, irreducible elliptic curves, C, of degree $d = 7, 8$, with $\mathcal{I}_C(4)$ globally generated, we have to recall some results due to Mori ([11]).

According to [11] Remark 4, Prop. 6, there exists a smooth quartic surface $S \subset \mathbb{P}^3$ such that $Pic(S) = \mathbb{Z}H \oplus \mathbb{Z}X$ where X is a smooth elliptic curve of degree d ($7 \leq d \leq 8$). The intersection pairing is given by: $H^2 = 4$, $X^2 = 0$, $H.X = d$. Such a surface doesn't contain any smooth rational curve ([11, p. 130]). In particular: $(*)$ every integral curve, Z, on S has degree ≥ 4 with equality if and only if Z is a planar quartic curve or an elliptic quartic curve.

Lemma 2.9. *With notations as above, $h^0(\mathcal{I}_X(3)) = 0$.*

Proof. A curve $Z \in |3H - X|$ has invariants $(d_Z, g_Z) = (5, -2)$ (if $d = 7$) or $(4, -5)$ (if $d = 8$), so Z is not integral. It follows that Z must contain an integral curve of degree < 4, but this is impossible. □

Lemma 2.10. *With notations as above $|4H - X|$ is base point free, hence there exist smooth, irreducible elliptic curves, X, of degree d, $7 \leq d \leq 8$, such that $\mathcal{I}_X(4)$ is globally generated.*

Proof. Let's first prove the following: *Claim:* Every curve in $|4H - X|$ is integral.

If $Y \in |4H - X|$ is not integral then $Y = Y_1 + Y_2$ where Y_1 is integral with $\deg(Y_1) = 4$ (observe that $\deg(Y) = 9$ or 8).

If Y_1 is planar then $Y_1 \sim H$, so $4H - X \sim H + Y_2$ and it follows that $3H \sim X + Y_2$, in contradiction with $h^0(\mathcal{I}_X(3)) = 0$ (cf 2.9).

So we may assume that Y_1 is a quartic elliptic curve, i.e. (i) $Y_1^2 = 0$ and (ii) $Y_1.H = 4$. Setting $Y_1 = aH + bX$, we get from (i): $2a(2a + bd) = 0$. Hence (α) $a = 0$, or (β) $2a + bd = 0$.

(α) In this case $Y_1 = bX$, hence (for degree reasons and since S doesn't contain curves of degree < 4), $Y_2 = \varnothing$ and $Y = X$, which is integral.

(β) Since $Y_1.H = 4$, we get $2a + (2a + bd) = 2a = 4$, hence $a = 2$ and $bd = -4$ which is impossible ($d = 7$ or 8 and $b \in \mathbb{Z}$).

This concludes the proof of the claim.

Since $(4H - X)^2 \geq 0$, the claim implies that $4H - X$ is numerically effective. Now we conclude by a result of Saint-Donat (cf. [11, Theorem 5]) that $|4H - X|$

is base point free, i.e. $\mathcal{I}_{X,S}(4)$ is globally generated. By the exact sequence: $0 \to \mathcal{O} \to \mathcal{I}_X(4) \to \mathcal{I}_{X,S}(4) \to 0$ we get that $\mathcal{I}_X(4)$ is globally generated. □

REMARK 2.11. *If $d = 8$, a general element $Y \in |4H - X|$ is a smooth elliptic curve of degree 8. By the way $Y \neq X$ (see [1]). The exact sequence of liaison: $0 \to \mathcal{I}_U(4) \to \mathcal{I}_X(4) \to \omega_Y \to 0$ shows that $h^0(\mathcal{I}_X(4)) = 3$ (i.e. X is of maximal rank). In case $d = 8$ Lemma 2.10 is stated in [2], however the proof there is incomplete, indeed in order to apply the enumerative formula of [8] one has to know that X is a* connected *component of $\bigcap_{i=1}^{3} F_i$; this amounts to say that the base locus of $|4H - X|$ on F_1 has dimension ≤ 0.*

To conclude we have:

LEMMA 2.12. *Let X be the disjoint union of two smooth, irreducible quartic elliptic cuvres, then $\mathcal{I}_X(4)$ is generated by global sections.*

Proof. Let $X = C_1 \sqcup C_2$. We have: $0 \to \mathcal{O}(-4) \to 2.\mathcal{O}(-2) \to \mathcal{I}_{C_1} \to 0$, twisting by $\mathcal{I}_{C_2}$, since $C_1 \cap C_2 = \varnothing$, we get:
$0 \to \mathcal{I}_{C_2}(-4) \to 2.\mathcal{I}_{C_2}(-2) \to \mathcal{I}_X \to 0$ and the result follows. □

Summarizing:

PROPOSITION 2.13. *There exists an indecomposable rank two vector bundle, E, on $\mathbb{P}^3$, generated by global sections and with $c_1(E) = 4$ if and only if $5 \leq c_2(E) \leq 8$ and in these cases there is an exact sequence:*

$$0 \to \mathcal{O} \to E \to \mathcal{I}_C(4) \to 0$$

where C is a smooth irreducible elliptic curve of degree $c_2(E)$ or, if $c_2(E) = 8$, the disjoint union of two smooth elliptic quartic curves.

2.4. Globally generated rank two vector bundles with $c_1 = 5$.

We start by listing the possible cases:

PROPOSITION 2.14. *If E is an indecomposable, globally generated, rank two vector bundle on $\mathbb{P}^3$ with $c_1(E) = 5$, then $c_2(E) \in \{8, 10, 12\}$ and there is an exact sequence:*

$$0 \to \mathcal{O} \to E \to \mathcal{I}_C(5) \to 0$$

where C is a smooth, irreducible curve of degree $d = c_2(E)$, with $\omega_C \simeq \mathcal{O}_C(1)$.

In any case E is stable.

Proof. A general section of E vanishes along a smooth curve, C, of degree $d = c_2(E)$ with $\omega_C \simeq \mathcal{O}_C(1)$. Hence every irreducible component, Y, of C is a smooth, irreducible curve with $\omega_Y \simeq \mathcal{O}_Y(1)$. In particular $\deg(Y) = 2g(Y) - 2$ is even and $\deg(Y) \geq 4$.

1. If $d = 4$, then C is a planar curve and E splits.
2. If $d = 6$, C is necessarily irreducible (of genus 4). It is well known that any such curve is a complete intersection $(2, 3)$, hence E splits.
3. If $d = 8$ and C is not irreducible, then $C = P_1 \sqcup P_2$, the disjoint union of two planar quartic curves. If $L = \langle P_1 \rangle \cap \langle P_2 \rangle$, then every quintic containing C contains L in contradiction with the fact that $\mathcal{I}_C(5)$ is generated by global sections. Hence C is irreducible.
4. If $d = 10$ and C is not irreducible, then $C = P \sqcup X$, where P is a planar curve of degree 4 and where X is a degree 6 curve (X is a complete intersection $(2, 3)$). Every quintic containing C vanishes on P and on the 8 points of $X \cap \langle P \rangle$, since these 8 points are not on a line, the quintic vanishes on the plane $\langle P \rangle$. This contradicts the fact that $\mathcal{I}_C(5)$ is globally generated.
5. If $d = 12$ and C is not irreducible we have three possibilities:
   (a) $C = P_1 \sqcup P_2 \sqcup P_3$, P_i planar quartic curves
   (b) $C = X_1 \sqcup X_2$, X_i complete intersection curves of types $(2, 3)$
   (c) $C = Y \sqcup P$, Y a canonical curve of degree 8, P a planar curve of degree 4.

   (a) This case is impossible (consider the line $\langle P_1 \rangle \cap \langle P_2 \rangle$).
   (b) We have $X_i = Q_i \cap F_i$. Let Z be the quartic curve $Q_1 \cap Q_2$. Then $X_i \cap Z = F_i \cap Z$, i.e. X_i meets Z in 12 points. It follows that every quintic containing C meets Z in 24 points, hence such a quintic contains Z. Again this contradicts the fact that $\mathcal{I}_C(5)$ is globally generated.
   (c) This case too is impossible: every quintic containing C vanishes on P and on the points $\langle P \rangle \cap Y$, hence on $\langle P \rangle$.

   We conclude that if $d = 12$, C is irreducible.

The normalized bundle is $E(-3)$, since in any case $h^0(\mathcal{I}_C(2)) = 0$ (every smooth irreducible subcanonical curve on a quadric surface is a complete intersection), E is stable. □

Now we turn to the existence part.

LEMMA 2.15. *There exist indecomposable rank two vector bundles on $\mathbb{P}^3$ with Chern classes $c_1 = 5$ and $c_2 \in \{8, 10\}$ which are globally generated.*

Proof. Let $R = \sqcup_{i=1}^{s} L_i$ be the union of s disjoint lines, $2 \leq s \leq 3$. We may perform a liaison $(s,3)$ and link R to $K = \sqcup_{i=1}^{s} K_i$, the union of s disjoint conics. The exact sequence of liaison: $0 \to \mathcal{I}_U(4) \to \mathcal{I}_K(4) \to \omega_R(5-s) \to 0$ shows that $\mathcal{I}_K(4)$ is globally generated (n.b. $5-s \geq 2$).
Since $\omega_K(1) \simeq \mathcal{O}_K$ we have an exact sequence: $0 \to \mathcal{O} \to \mathcal{E}(2) \to \mathcal{I}_K(3) \to 0$, where $\mathcal{E}$ is a rank two vector bundle with Chern classes $c_1 = -1, c_2 = 2s-2$. Twisting by $\mathcal{O}(1)$ we get: $0 \to \mathcal{O}(1) \to \mathcal{E}(3) \to \mathcal{I}_K(4) \to 0$ $(*)$. The Chern classes of $\mathcal{E}(3)$ are $c_1 = 5$, $c_2 = 2s+4$ (i.e. $c_2 = 8, 10$). Since $\mathcal{I}_K(4)$ is globally generated, it follows from $(*)$ that $\mathcal{E}(3)$ too, is generated by global sections. □

Remark 2.16.

1. *If $\mathcal{E}$ is as in the proof of Lemma 2.15 a general section of $\mathcal{E}(3)$ vanishes along a smooth, irreducible (because $h^1(\mathcal{E}(-2)) = 0$) canonical curve, C, of genus $1 + c_2/2$ ($g = 5, 6$) such that $\mathcal{I}_C(5)$ is globally generated. By construction these curves are not of maximal rank ($h^0(\mathcal{I}_C(3)) = 1$ if $g = 5$, $h^0(\mathcal{I}_C(4)) = 2$ if $g = 6$). As explained in [9] 4 this is a general fact: no canonical curve of genus $g, 5 \leq g \leq 6$ in $\mathbb{P}^3$ is of maximal rank. We don't know if this is still true for $g = 7$.*

2. *According to [9] the general canonical curve of genus 6 lies on a unique quartic surface.*

3. *The proof of 2.15 breaks down with four conics: $\mathcal{I}_K(4)$ is no longer globally generated, every quartic containing K vanishes along the lines L_i ($5 - s = 1$). Observe also that four disjoint lines always have a quadrisecant and hence are an exception to the* normal generation conjecture*(the omogeneous ideal is not generated in degree three as it should be).*

Remark 2.17. *The case $(c_1, c_2) = (5, 12)$ remains open. It can be shown that if E exists, a general section of E is linked, by a complete intersections of two quintics, to a smooth, irreducible curve, X, of degree 13, genus 10 having $\omega_X(-1)$ as a base point free g^1_5. One can prove the existence of curves $X \subset \mathbb{P}^3$, smooth, irreducible, of degree 13, genus 10, with $\omega_X(-1)$ a base point free pencil and lying on* one *quintic surface. But we are unable to show the existence of such a curve with $h^0(\mathcal{I}_X(5)) \geq 3$ (or even with $h^0(\mathcal{I}_X(5)) \geq 2$). We believe that such bundles do not exist.*

3. Globally generated rank two vector bundles on $\mathbb{P}^n$, $n \geq 4$.

For $n \geq 4$ and $c_1 \leq 5$ there is no surprise:

Proposition 3.1. *Let E be a globally generated rank two vector bundle on $\mathbb{P}^n$, $n \geq 4$. If $c_1(E) \leq 5$, then E splits.*

Proof. It is enough to treat the case $n = 4$. A general section of E vanishes along a smooth (irreducible) subcanonical surface, S: $0 \to \mathcal{O} \to E \to \mathcal{I}_S(c_1) \to 0$. By [5], if $c_1 \leq 4$, then S is a complete intersection and E splits. Assume now $c_1 = 5$. Consider the restriction of E to a general hyperplane H. If E doesn't split, by 2.14 we get that the normalized Chern classes of E are: $c_1 = -1$, $c_2 \in \{2, 4, 6\}$. By Schwarzenberger condition: $c_2(c_2 + 2) \equiv 0$ $(mod\ 12)$. The only possibilities are $c_2 = 4$ or $c_2 = 6$. If $c_2 = 4$, since E is stable (because $E|H$ is, see 2.14), we have ([3]) that E is a Horrocks-Mumford bundle. But the Horrocks-Mumford bundle (with $c_1 = 5$) is not globally generated.

The case $c_2 = 6$ is impossible: such a bundle would yield a smooth surface $S \subset \mathbb{P}^4$, of degree 12 with $\omega_S \simeq \mathcal{O}_S$, but the only smooth surface with $\omega_S \simeq \mathcal{O}_S$ in $\mathbb{P}^4$ is the abelian surface of degree 10 of Horrocks-Mumford. □

Remark 3.2. *For $n > 4$ the results in [6] give stronger and stronger (as n increases) conditions for the existence of indecomposable rank two vector bundles generated by global sections.*

Putting everything together, the proof of Theorem 1.1 is complete.

References

[1] V. Beorchia and Ph. Ellia, *Normal bundle and complete intersections*, Rend. Sem. Mat. Univ. Politec. Torino **48** (1990), 553–562.

[2] J. D'Almeida, *Une involution sur un espace de modules de fibrés instantons*, Bull. Soc. Math. France **128** (2000), 577–584.

[3] W. Decker, *Stable rank 2 vector bundles with Chern classes $c_1 = -1, c_2 = 4$*, Math. Ann. **275** (1986), 481–500.

[4] Ph. Ellia, *Sur les fibrés uniformes de rang $n + 1$ sur $\mathbb{P}^n$*, Mém. Soc. Math. France **7** (1982).

[5] Ph. Ellia, D. Franco, and L. Gruson, *On subcanonical surfaces of $\mathbb{P}^4$*, Math. Z. **251** (2005), 257–265.

[6] Ph. Ellia, D. Franco, and L. Gruson, *Smooth divisors of projective hypersurfaces*, Comment. Math. Helv. **83** (2008), 371–385.

[7] M.L. Fania and E. Mezzetti, *Vector spaces of skew-symmetric matrices of constant rank*, Linear Algebra Appl. **434** (2011), 2383–2403.

[8] W. Fulton, *Intersection theory*, Ergeb. Math. Grenzgeb., no. 2, Springer, Berlin, 1984.

[9] L. Gruson and Ch. Peskine, *Genre des courbes de l'espace projectif*, Lecture Notes in Math. **687** (1978), 31–59.

[10] S. Huh, *On triple Veronese embeddings of $\mathbb{P}^n$ in the Grassmannians*, Math. Nachr. **284** (2011), 1453–1461.

[11] S. Mori, *On degrees and genera of curves on smooth quartic surfaces in $\mathbb{P}^3$*, Nagoya Math. J. **96** (1984), 127–132.

[12] J.C. Sierra and L. Ugaglia, *On double Veronese embeddings in the Grassmannian $G(1, N)$*, Math. Nachr. **279** (2006), 798–804.

[13] J.C. Sierra and L. Ugaglia, *On globally generated vector bundles on projective spaces*, J. Pure Appl. Algebra **213** (2009), 2141–2146.
[14] A. Van de Ven, *On uniform vector bundles*, Math. Ann. **195** (1972), 245–248.

Authors' addresses:

Ludovica Chiodera
Dipartimento di Matematica
Università di Ferrara
via Machiavelli 35, 44100 Ferrara, Italy
E-mail: `ludovica.chiodera@unife.it`

Philippe Ellia
Dipartimento di Matematica
Università di Ferrara
via Machiavelli 35, 44100 Ferrara, Italy
E-mail: `phe@unife.it`

Received November 30, 2011
Revised April 2, 2012