

# Weakly $\omega b$ -Continuous Functions<sup>1</sup>

JAMAL M. MUSTAFA

ABSTRACT. *In this paper we introduce a new class of functions called weakly  $\omega b$ -continuous functions and investigate several properties and characterizations. Connections with other existing concepts, such as  $\omega b$ -continuous and weakly  $b$ -continuous functions, are also discussed.*

Keywords:  $b$ -Open Sets,  $\omega b$ -Open Sets, Weakly  $b$ -Continuous Functions, Weakly  $\omega b$ -Continuous Functions.

MS Classification 2010: 54C05, 54C08

## 1. Introduction

The notion of  $b$ -open sets in topological spaces was introduced in 1996 by Andrijevic [1]. This type of sets discussed by El-Atik [2] under the name of  $\gamma$ -open sets. In 2008, Noiri, Al-Omari and Noorani [4] introduced the notions of  $\omega b$ -open sets and  $\omega b$ -continuous functions. We continue to introduce and study properties and characterizations of weakly  $\omega b$ -continuous functions.

Let  $A$  be a subset of a space  $(X, \tau)$ . The closure ( resp. interior ) of  $A$  will be denoted by  $Cl(A)$  ( resp.  $Int(A)$  ).

A subset  $A$  of a space  $(X, \tau)$  is called  $b$ -open [1] if  $A \subseteq Cl(Int(A)) \cup Int(Cl(A))$ . The complement of a  $b$ -open set is called a  $b$ -closed set. The union of all  $b$ -open sets contained in  $A$  is called the  $b$ -interior of  $A$ , denoted by  $bInt(A)$  and the intersection of all  $b$ -closed sets containing  $A$  is called the  $b$ -closure of  $A$ , denoted by  $bCl(A)$ . The family of all  $b$ -open ( resp.  $b$ -closed ) sets in  $(X, \tau)$  is denoted by  $BO(X)$  ( resp.  $BC(X)$  ).

DEFINITION 1.1. *A subset  $A$  of a space  $X$  is said to be  $\omega b$ -open [4] if for every  $x \in A$ , there exists a  $b$ -open subset  $U_x \subseteq X$  containing  $x$  such that  $U_x - A$  is countable.*

The complement of an  $\omega b$ -open set is said to be  $\omega b$ -closed [4]. The intersection of all  $\omega b$ -closed sets of  $X$  containing  $A$  is called the  $\omega b$ -closure of  $A$  and is denoted by  $\omega bCl(A)$ . The union of all  $\omega b$ -open sets of  $X$  contained in  $A$  is called the  $\omega b$ -interior of  $A$  and is denoted by  $\omega bInt(A)$ .

---

<sup>1</sup>Jamal M. Mustafa did this research during the sabbatical leave from Department of Mathematics, Al al-Bayt University, Mafraq, Jordan.

LEMMA 1.2 ([4]). For a subset of a topological space,  $b$ -openness implies  $\omega b$  – openness.

LEMMA 1.3 ([4]). The intersection of an  $\omega b$  – open set with an open set is  $\omega b$  – open.

LEMMA 1.4 ([4]). The union of any family of  $\omega b$  – open sets is  $\omega b$  – open.

## 2. Weakly $\omega b$ -Continuous Functions

DEFINITION 2.1. A function  $f : (X, \tau) \rightarrow (Y, \rho)$  is said to be:

- (a)  $\omega b$ -continuous [4] if for each  $x \in X$  and each open set  $V$  in  $Y$  containing  $f(x)$ , there exists an  $\omega b$  – open set  $U$  in  $X$  containing  $x$  such that  $f(U) \subseteq V$ .
- (b) weakly  $b$ -continuous [7] if for each  $x \in X$  and each open set  $V$  in  $Y$  containing  $f(x)$ , there exists a  $b$ -open set  $U$  in  $X$  containing  $x$  such that  $f(U) \subseteq Cl(V)$ .

DEFINITION 2.2. A function  $f : (X, \tau) \rightarrow (Y, \rho)$  is said to be weakly  $\omega b$ -continuous if for each  $x \in X$  and each open set  $V$  in  $Y$  containing  $f(x)$ , there exists an  $\omega b$  – open set  $U$  in  $X$  containing  $x$  such that  $f(U) \subseteq Cl(V)$ .

REMARK 2.3. Every  $\omega b$ -continuous function is weakly  $\omega b$ -continuous, but the converse is not true in general as the following example shows.

EXAMPLE 2.4. Let  $X = R$  with the usual topology  $\tau$  and  $Y = \{a, b\}$  with  $\rho = \{\phi, Y, \{a\}\}$ . Define a function  $f : (X, \tau) \rightarrow (Y, \rho)$  by  $f(x) = a$  if  $x \in Q$  and  $f(x) = b$  if  $x \in R - Q$ . Then  $f$  is weakly  $\omega b$ -continuous but not  $\omega b$ -continuous.

REMARK 2.5. Since every  $b$  – open set is  $\omega b$  – open then every weakly  $b$ -continuous function is weakly  $\omega b$ -continuous but the converse is not true in general as the following example shows.

EXAMPLE 2.6. Let  $X = Y = \{a, b, c\}$ ,  $\tau = \{\phi, X, \{c\}, \{a, b\}, \{a, b, c\}\}$  and  $\rho = \{\phi, Y, \{a, b\}, \{c, d\}\}$ . Define a function  $f : (X, \tau) \rightarrow (Y, \rho)$  by  $f(a) = a$ ,  $f(b) = d$ ,  $f(c) = c$  and  $f(d) = b$ . Then  $f$  is weakly  $\omega b$ -continuous but not weakly  $b$ -continuous.

THEOREM 2.7. A function  $f : (X, \tau) \rightarrow (Y, \rho)$  is weakly  $\omega b$ -continuous if and only if for every open set  $V$  in  $Y$ ,  $f^{-1}(V) \subseteq \omega bInt[f^{-1}(Cl(V))]$ .

*Proof.*

- $\Rightarrow$ ) Let  $V \in \rho$  and  $x \in f^{-1}(V)$ . Then there exists an  $\omega b$  – open set  $U$  in  $X$  such that  $x \in U$  and  $f(U) \subseteq Cl(V)$ . Therefore, we have  $x \in U \subseteq f^{-1}(Cl(V))$  and hence  $x \in \omega bInt[f^{-1}(Cl(V))]$  which means that  $f^{-1}(V) \subseteq \omega bInt[f^{-1}(Cl(V))]$ .

$\Leftarrow$ ) Let  $x \in X$  and  $V \in \rho$  with  $f(x) \in V$ . Then  $x \in f^{-1}(V) \subseteq \omega bInt[f^{-1}(Cl(V))]$ . Let  $U = \omega bInt[f^{-1}(Cl(V))]$ . Then  $U$  is  $\omega b$ -open and  $f(U) \subseteq Cl(V)$ .  $\square$

**THEOREM 2.8.** *Let  $f : (X, \tau) \rightarrow (Y, \rho)$  be a weakly  $\omega b$ -continuous function. If  $V$  is a clopen subset of  $Y$ , then  $f^{-1}(V)$  is  $\omega b$ -open and  $\omega b$ -closed in  $X$ .*

*Proof.* Let  $x \in X$  and  $V$  be a clopen subset of  $Y$  such that  $f(x) \in V$ . Then there exists an  $\omega b$ -open set  $U$  in  $X$  containing  $x$  such that  $f(U) \subseteq Cl(V)$ . Hence  $x \in U$  and  $f(U) \subseteq V$  and so  $x \in U \subseteq f^{-1}(V)$ . This shows that  $f^{-1}(V)$  is  $\omega b$ -open in  $X$ . Since  $Y - V$  is a clopen set in  $Y$ , so  $f^{-1}(Y - V)$  is  $\omega b$ -open in  $X$ . But  $f^{-1}(Y - V) = X - f^{-1}(V)$ . Therefore  $f^{-1}(V)$  is  $\omega b$ -closed in  $X$ . Hence  $f^{-1}(V)$  is  $\omega b$ -open and  $\omega b$ -closed in  $X$ .  $\square$

**THEOREM 2.9.** *A function  $f : (X, \tau) \rightarrow (Y, \rho)$  is weakly  $\omega b$ -continuous if and only if for every closed set  $C$  in  $Y$ ,  $\omega bCl[f^{-1}(Int(C))] \subseteq f^{-1}(C)$ .*

*Proof.*

$\Rightarrow$ ) Let  $C$  be a closed set in  $Y$ . Then  $Y - C$  is an open set in  $Y$  so by Theorem 2.8  $f^{-1}(Y - C) \subseteq \omega bInt[f^{-1}(Cl(Y - C))] = \omega bInt[f^{-1}(Y - Int(C))] = X - \omega bCl[f^{-1}(Int(C))]$ . Thus  $\omega bCl[f^{-1}(Int(C))] \subseteq f^{-1}(C)$ .

$\Leftarrow$ ) Let  $x \in X$  and  $V \in \rho$  with  $f(x) \in V$ . So  $Y - V$  is a closed set in  $Y$ . So by assumption  $\omega bCl[f^{-1}(Int(Y - V))] \subseteq f^{-1}(Y - V)$ . Thus  $x \notin \omega bCl[f^{-1}(Int(Y - V))]$ . Hence there exists an  $\omega b$ -open set  $U$  in  $X$  such that  $x \in U$  and  $U \cap f^{-1}(Int(Y - V)) = \phi$  which implies that  $f(U) \cap Int(Y - V) = \phi$ . Then  $f(U) \subseteq Y - Int(Y - V)$ , so  $f(U) \subseteq Cl(V)$ , which means that  $f$  is weakly  $\omega b$ -continuous.  $\square$

**THEOREM 2.10.** *Let  $f : (X, \tau) \rightarrow (Y, \rho)$  be a surjection function such that  $f(U)$  is  $\omega b$ -open in  $Y$  for any  $\omega b$ -open set  $U$  in  $X$  and let  $g : (Y, \rho) \rightarrow (Z, \sigma)$  be any function. If  $g \circ f$  is weakly  $\omega b$ -continuous then  $g$  is weakly  $\omega b$ -continuous.*

*Proof.* Let  $y \in Y$ . Since  $f$  is surjection, there exists  $x \in X$  such that  $f(x) = y$ . Let  $V \in \sigma$  with  $g(y) \in V$ , so  $(g \circ f)(x) \in V$ . Since  $g \circ f$  is weakly  $\omega b$ -continuous there exists an  $\omega b$ -open set  $U$  in  $X$  containing  $x$  such that  $(g \circ f)(U) \subseteq Cl(V)$ . By assumption  $H = f(U)$  is an  $\omega b$ -open set in  $Y$  and contains  $f(x) = y$ . Thus  $g(H) \subseteq Cl(V)$ . Hence  $g$  is weakly  $\omega b$ -continuous.  $\square$

**DEFINITION 2.11.** *A function  $f : (X, \tau) \rightarrow (Y, \rho)$  is called  $\omega b$ -irresolute if  $f^{-1}(V)$  is  $\omega b$ -open in  $(X, \tau)$  for every  $\omega b$ -open set  $V$  in  $(Y, \rho)$ .*

**THEOREM 2.12.** *If  $f : (X, \tau) \rightarrow (Y, \rho)$  is  $\omega b$ -irresolute and  $g : (Y, \rho) \rightarrow (Z, \sigma)$  is weakly  $\omega b$ -continuous then  $g \circ f : (X, \tau) \rightarrow (Z, \sigma)$  is weakly  $\omega b$ -continuous.*

*Proof.* Let  $x \in X$  and  $V \in \sigma$  such that  $(gof)(x) = g(f(x)) \in V$ . Let  $y = f(x)$ . Since  $g$  is weakly  $\omega b$ -continuous. So there exists an  $\omega b$ -open set  $W$  in  $Y$  such that  $y \in W$  and  $g(W) \subseteq Cl(V)$ . Let  $U = f^{-1}(W)$ . Then  $U$  is an  $\omega b$ -open set in  $X$  as  $f$  is  $\omega b$ -irresolute. Now  $(gof)(U) = g(f(f^{-1}(W))) \subseteq g(W)$ . Then  $x \in U$  and  $(gof)(U) \subseteq Cl(V)$ . Hence  $gof$  is weakly  $\omega b$ -continuous.  $\square$

**THEOREM 2.13.** *If  $f : (X, \tau) \rightarrow (Y, \rho)$  is weakly  $\omega b$ -continuous and  $g : (Y, \rho) \rightarrow (Z, \sigma)$  is continuous then  $gof : (X, \tau) \rightarrow (Z, \sigma)$  is weakly  $\omega b$ -continuous.*

*Proof.* Let  $x \in X$  and  $W$  be an open set in  $Z$  containing  $(gof)(x) = g(f(x))$ . Then  $g^{-1}(W)$  is an open set in  $Y$  containing  $f(x)$ . So there exists an  $\omega b$ -open set  $U$  in  $X$  containing  $x$  such that  $f(U) \subseteq Cl(g^{-1}(W))$ . Since  $g$  is continuous we have  $(gof)(U) \subseteq g(Cl(g^{-1}(W))) \subseteq g(g^{-1}(Cl(W))) \subseteq Cl(W)$ .  $\square$

**THEOREM 2.14.** *A function  $f : X \rightarrow Y$  is weakly  $\omega b$ -continuous if and only if the graph function  $g : X \rightarrow X \times Y$  of  $f$  defined by  $g(x) = (x, f(x))$  for each  $x \in X$ , is weakly  $\omega b$ -continuous.*

*Proof.*

$\Rightarrow$ ) Suppose that  $f$  is weakly  $\omega b$ -continuous. Let  $x \in X$  and  $W$  be an open set in  $X \times Y$  containing  $g(x)$ . Then there exists a basic open set  $U_1 \times V$  in  $X \times Y$  such that  $g(x) = (x, f(x)) \in U_1 \times V \subseteq W$ . Since  $f$  is weakly  $\omega b$ -continuous there exists an  $\omega b$ -open set  $U_2$  in  $X$  containing  $x$  such that  $f(U_2) \subseteq Cl(V)$ . Let  $U = U_1 \cap U_2$  then  $U$  is an  $\omega b$ -open set in  $X$  with  $x \in U$  and  $g(U) \subseteq Cl(W)$ .

$\Leftarrow$ ) Suppose that  $g$  is weakly  $\omega b$ -continuous. Let  $x \in X$  and  $V$  be an open set in  $Y$  containing  $f(x)$ . Then  $X \times V$  is an open set containing  $g(x)$  and hence there exists an  $\omega b$ -open set  $U$  in  $X$  containing  $x$  such that  $g(U) \subseteq Cl(X \times V) = X \times Cl(V)$ . Therefore, we have  $f(U) \subseteq Cl(V)$  and hence  $f$  is weakly  $\omega b$ -continuous.  $\square$

**THEOREM 2.15.** *If  $f : (X, \tau) \rightarrow (Y, \rho)$  is a weakly  $\omega b$ -continuous function and  $Y$  is Hausdorff then the set  $G(f) = \{(x, f(x)) : x \in X\}$  is an  $\omega b$ -closed set in  $X \times Y$ .*

*Proof.* Let  $(x, y) \in (X \times Y) - G(f)$ . Then  $y \neq f(x)$ . Since  $Y$  is Hausdorff, there exist two disjoint open sets  $U$  and  $V$  such that  $y \in U$  and  $f(x) \in V$ . Since  $f$  is weakly  $\omega b$ -continuous, there exists an  $\omega b$ -open set  $W$  containing  $x$  such that  $f(W) \subseteq Cl(V)$ . Since  $V$  and  $U$  are disjoint, we have  $U \cap Cl(V) = \phi$  and hence  $U \cap f(W) = \phi$ . This shows that  $(W \times U) \cap G(f) = \phi$ . Then  $G(f)$  is  $\omega b$ -closed.  $\square$

**THEOREM 2.16.** *If  $f : X_1 \rightarrow Y$  is  $\omega b$ -continuous,  $g : X_2 \rightarrow Y$  is weakly  $\omega b$ -continuous and  $Y$  is Hausdorff, then the set  $A = \{(x_1, x_2) \in X_1 \times X_2 : f(x_1) = g(x_2)\}$  is  $\omega b$ -closed in  $X_1 \times X_2$ .*

*Proof.* Let  $(x_1, x_2) \in (X_1 \times X_2) - A$ . Then  $f(x_1) \neq g(x_2)$  and there exist open sets  $V_1$  and  $V_2$  in  $Y$  such that  $f(x_1) \in V_1$ ,  $g(x_2) \in V_2$  and  $V_1 \cap V_2 = \phi$ , hence  $V_1 \cap Cl(V_2) = \phi$ . Since  $f$  is  $\omega b$ -continuous there exists an  $\omega b$ -open set  $U_1$  in  $X_1$  containing  $x_1$  such that  $f(U_1) \subseteq V_1$ . Since  $g$  is weakly  $\omega b$ -continuous there exists an  $\omega b$ -open set  $U_2$  in  $X_2$  containing  $x_2$  such that  $g(U_2) \subseteq Cl(V_2)$ . Now  $U_1 \times U_2$  is an  $\omega b$ -open set in  $X_1 \times X_2$  with  $(x_1, x_2) \in U_1 \times U_2 \subseteq (X_1 \times X_2) - A$ . This shows that  $A$  is  $\omega b$ -closed in  $X_1 \times X_2$ .  $\square$

**THEOREM 2.17.** *If  $(Y, \rho)$  is a regular space then a function  $f : (X, \tau) \rightarrow (Y, \rho)$  is weakly  $\omega b$ -continuous if and only if it is  $\omega b$ -continuous*

*Proof.*

$\Rightarrow$ ) Let  $x$  be any point in  $X$  and  $V$  be any open set in  $Y$  containing  $f(x)$ . Since  $(Y, \rho)$  is regular, there exists  $W \in \rho$  such that  $f(x) \in W \subseteq Cl(W) \subseteq V$ . Since  $f$  is weakly  $\omega b$ -continuous there exists an  $\omega b$ -open set  $U$  in  $X$  containing  $x$  such that  $f(U) \subseteq Cl(W)$ . So  $f(U) \subseteq V$ . Therefore,  $f$  is  $\omega b$ -continuous.

$\Leftarrow$ ) Clear.  $\square$

**DEFINITION 2.18.** *Any weakly  $\omega b$ -continuous function  $f : X \rightarrow A$ , where  $A \subseteq X$  and  $f_A = f|_A$  is the identity function on  $A$ , is called weakly  $\omega b$ -continuous retraction.*

**THEOREM 2.19.** *Let  $f : X \rightarrow A$  be a weakly  $\omega b$ -continuous retraction of  $X$  onto  $A$  where  $A \subseteq X$ . If  $X$  is a Hausdorff space, then  $A$  is an  $\omega b$ -closed set in  $X$ .*

*Proof.* Suppose that  $A$  is not  $\omega b$ -closed in  $X$ . Then there exist a point  $x \in \omega bCl(A) - A$ . Since  $f$  is weakly  $\omega b$ -continuous retraction, we have  $f(x) \neq x$ . Since  $X$  is Hausdorff, there exist two disjoint open sets  $U$  and  $V$  such that  $x \in U$  and  $f(x) \in V$ . Then we have  $U \cap Cl(V) = \phi$ . Now, Let  $W$  be any  $\omega b$ -open set in  $X$  containing  $x$ . Then  $U \cap W$  is an  $\omega b$ -open set containing  $x$  and hence  $(U \cap W) \cap A \neq \phi$  because  $x \in \omega bCl(A)$ . Let  $y \in (U \cap W) \cap A$ . Since  $y \in A$ ,  $f(y) = y \in U$  and hence  $f(y) \notin Cl(V)$ . This gives that  $f(W)$  is not a subset of  $Cl(V)$ . This contradicts the fact that  $f$  is weakly  $\omega b$ -continuous. Therefore  $A$  is  $\omega b$ -closed in  $X$ .  $\square$

**DEFINITION 2.20.** *A space  $X$  is called:*

- (a)  $\omega b - T_1$  if for each pair of distinct points  $x$  and  $y$  in  $X$ , there exist two  $\omega b$ -open sets  $U$  and  $V$  of  $X$  containing  $x$  and  $y$ , respectively, such that  $y \notin U$  and  $x \notin V$ .
- (b)  $\omega b - T_2$  if for each pair of distinct points  $x$  and  $y$  in  $X$ , there exist two  $\omega b$ -open sets  $U$  and  $V$  of  $X$  containing  $x$  and  $y$ , respectively, such that  $U \cap V = \phi$

**THEOREM 2.21.** *If for each pair of distinct points  $x$  and  $y$  in a space  $X$  there exists a function  $f$  of  $X$  into a Hausdorff space  $Y$  such that*

- 1)  $f(x) \neq f(y)$
- 2)  $f$  is  $\omega b$ -continuous at  $x$  and
- 3)  $f$  is weakly  $\omega b$ -continuous at  $y$ ,

*then  $X$  is  $\omega b - T_2$ .*

*Proof.* Since  $f(x) \neq f(y)$  and  $Y$  is Hausdorff, there exist open sets  $V_1$  and  $V_2$  of  $Y$  containing  $f(x)$  and  $f(y)$ , respectively, such that  $V_1 \cap V_2 = \phi$ , hence  $V_1 \cap Cl(V_2) = \phi$ . Since  $f$  is  $\omega b$ -continuous at  $x$ , there exists an  $\omega b$ -open set  $U_1$  in  $X$  containing  $x$  such that  $f(U_1) \subseteq V_1$ . Since  $f$  is weakly  $\omega b$ -continuous at  $y$ , there exists an  $\omega b$ -open set  $U_2$  in  $X$  containing  $y$  such that  $f(U_2) \subseteq Cl(V_2)$ . Therefore we obtain  $U_1 \cap U_2 = \phi$ . This shows that  $X$  is  $\omega b - T_2$ .  $\square$

**DEFINITION 2.22.** *A space  $X$  is called Urysohn [5] if for each pair of distinct points  $x$  and  $y$  in  $X$ , there exist open sets  $U$  and  $V$  such that  $x \in U$ ,  $y \in V$  and  $Cl(U) \cap Cl(V) = \phi$ .*

**THEOREM 2.23.** *Let  $f : (X, \tau) \rightarrow (Y, \rho)$  be a weakly  $\omega b$ -continuous injection. Then the following hold:*

- (a) *If  $Y$  is Hausdorff, then  $X$  is  $\omega b - T_1$ .*
- (b) *If  $Y$  is Urysohn, then  $X$  is  $\omega b - T_2$ .*

*Proof.*

- (a) Let  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ . Then  $f(x_1) \neq f(x_2)$  and there exist open sets  $V_1$  and  $V_2$  in  $Y$  containing  $f(x_1)$  and  $f(x_2)$ , respectively, such that  $V_1 \cap V_2 = \phi$ . Then we obtain  $f(x_1) \notin Cl(V_2)$  and  $f(x_2) \notin Cl(V_1)$ . Since  $f$  is weakly  $\omega b$ -continuous, there exist  $\omega b$ -open sets  $U_1$  and  $U_2$  with  $x_1 \in U_1$  and  $x_2 \in U_2$  such that  $f(U_1) \subseteq Cl(V_1)$  and  $f(U_2) \subseteq Cl(V_2)$ . Hence we obtain  $x_2 \notin U_1$  and  $x_1 \notin U_2$ . This shows that  $X$  is  $\omega b - T_1$ .
- (b) Let  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ . Then  $f(x_1) \neq f(x_2)$  and there exist open sets  $V_1$  and  $V_2$  in  $Y$  containing  $f(x_1)$  and  $f(x_2)$ , respectively, such that  $Cl(V_1) \cap Cl(V_2) = \phi$ . Since  $f$  is weakly  $\omega b$ -continuous there exist  $\omega b$ -open sets  $U_1$  and  $U_2$  in  $X$  with  $x_1 \in U_1$  and  $x_2 \in U_2$  such that  $f(U_1) \subseteq Cl(V_1)$  and  $f(U_2) \subseteq Cl(V_2)$ . Since  $f^{-1}(Cl(V_1)) \cap f^{-1}(Cl(V_2)) = \phi$  we obtain  $U_1 \cap U_2 = \phi$ . Hence  $X$  is  $\omega b - T_2$ .  $\square$

**DEFINITION 2.24.** *A function  $f : X \rightarrow Y$  is said to have a strongly  $\omega b$ -closed graph if for each  $(x, y) \in (X \times Y) - G(f)$  there exist an  $\omega b$ -open subset  $U$  of  $X$  and an open subset  $V$  of  $Y$  such that  $(x, y) \in U \times V$  and  $(U \times Cl(V)) \cap G(f) = \phi$ .*

**THEOREM 2.25.** *If  $Y$  is a Urysohn space and  $f : X \rightarrow Y$  is weakly  $\omega b$ -continuous, then  $G(f)$  is strongly  $\omega b$ -closed.*

*Proof.* Let  $(x, y) \in (X \times Y) - G(f)$ . Then  $y \neq f(x)$  and there exist open sets  $V$  and  $W$  in  $Y$  with  $f(x) \in V$  and  $y \in W$  such that  $Cl(V) \cap Cl(W) = \phi$ . Since  $f$  is weakly  $\omega b$ -continuous, there exists an  $\omega b$ -open subset  $U$  of  $X$  containing  $x$  such that  $f(U) \subseteq Cl(V)$ . Therefore we obtain  $f(U) \cap Cl(W) = \phi$  and hence  $(U \times Cl(W)) \cap G(f) = \phi$ . This shows that  $G(f)$  is strongly  $\omega b$ -closed in  $X \times Y$ .  $\square$

**THEOREM 2.26.** *Let  $f : (X, \tau) \rightarrow (Y, \rho)$  be a weakly  $\omega b$ -continuous function having strongly  $\omega b$ -closed graph  $G(f)$ . If  $f$  is injective, then  $X$  is  $\omega b - T_2$ .*

*Proof.* Let  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ . Since  $f$  is injective,  $f(x_1) \neq f(x_2)$  and  $(x_1, f(x_2)) \notin G(f)$ . Since  $G(f)$  is strongly  $\omega b$ -closed, there exist an  $\omega b$ -open subset  $U$  of  $X$  containing  $x_1$  and an open subset  $V$  of  $Y$  such that  $(x_1, f(x_2)) \in U \times V$  and  $(U \times Cl(V)) \cap G(f) = \phi$  and hence  $f(U) \cap Cl(V) = \phi$ . Since  $f$  is weakly  $\omega b$ -continuous, there exists an  $\omega b$ -open subset  $W$  of  $X$  containing  $x_2$  such that  $f(W) \subseteq Cl(V)$ . Therefore, we have  $f(U) \cap f(W) = \phi$  and hence  $U \cap W = \phi$ . This shows that  $X$  is  $\omega b - T_2$ .  $\square$

**DEFINITION 2.27.** *A space  $X$  is said to be  $\omega b$ -connected if  $X$  can not be written as a union of two non-empty disjoint  $\omega b$ -open sets.*

**THEOREM 2.28.** *If  $X$  is an  $\omega b$ -connected space and  $f : X \rightarrow Y$  is weakly  $\omega b$ -continuous surjection then  $Y$  is connected.*

*Proof.* Suppose that  $Y$  is not connected. Then there exist two non-empty disjoint open sets  $U$  and  $V$  in  $Y$  such that  $U \cup V = Y$ . Hence, we have  $f^{-1}(U) \cap f^{-1}(V) = \phi$ ,  $f^{-1}(U) \cup f^{-1}(V) = X$  and since  $f$  is surjection we have  $f^{-1}(U) \neq \phi \neq f^{-1}(V)$ . By Theorem 2.8, we have  $f^{-1}(U) \subseteq \omega bInt[f^{-1}(Cl(U))]$  and  $f^{-1}(V) \subseteq \omega bInt[f^{-1}(Cl(V))]$ . Since  $U$  and  $V$  are clopen we have  $f^{-1}(U) \subseteq \omega bInt[f^{-1}(U)]$  and  $f^{-1}(V) \subseteq \omega bInt[f^{-1}(V)]$  and hence  $f^{-1}(U)$  and  $f^{-1}(V)$  are  $\omega b$ -open. This implies that  $X$  is not  $\omega b$ -connected which is a contradiction. Therefore  $Y$  is connected.  $\square$

**DEFINITION 2.29.** *A topological space  $(X, \tau)$  is said to be:*

- (a) *almost compact [3] if every open cover of  $X$  has a finite subfamily whose closures cover  $X$ .*
- (b) *almost Lindelöf [6] if every open cover of  $X$  has a countable subfamily whose closures cover  $X$ .*

**DEFINITION 2.30.** *A topological space  $(X, \tau)$  is said to be  $\omega b$ -compact (resp.  $\omega b$ -Lindelöf) if every  $\omega b$ -open cover of  $X$  has a finite (resp. countable) subcover.*

**THEOREM 2.31.** *Let  $f : X \rightarrow Y$  be a weakly  $\omega b$ -continuous surjection. Then the following hold:*

- (a) *If  $X$  is  $\omega b$ -compact, then  $Y$  is almost compact.*
- (b) *If  $X$  is  $\omega b$ -Lindelöf, then  $Y$  is almost Lindelöf.*

*Proof.*

- (a) Let  $\{V_\alpha : \alpha \in \Delta\}$  be a cover of  $Y$  by open sets in  $Y$ . For each  $x \in X$  there exists  $V_{\alpha_x} \in \{V_\alpha : \alpha \in \Delta\}$  such that  $f(x) \in V_{\alpha_x}$ . Since  $f$  is weakly  $\omega b$ -continuous, there exists an  $\omega b$ -open set  $U_x$  of  $X$  containing  $x$  such that  $f(U_x) \subseteq Cl(V_{\alpha_x})$ . The family  $\{U_x : x \in X\}$  is a cover of  $X$  by  $\omega b$ -open sets of  $X$  and hence there exists a finite subset  $X_0$  of  $X$  such that  $X \subseteq \cup\{U_x : x \in X_0\}$ . Therefore, we obtain  $Y = f(X) \subseteq \cup\{Cl(V_{\alpha_x}) : x \in X_0\}$ . This shows that  $Y$  is almost compact.

- (b) Similar to (a). □

#### REFERENCES

- [1] D. ANDRIJEVIĆ, *On  $b$ -open sets*, Mat. Vesnik **48** (1996), 59–64.
- [2] A.A. EL-ATIK, *A study on some types of mappings on topological spaces*, M.Sc. Thesis, Tanta University, Egypt (1997).
- [3] T. NOIRI, *Between continuity and weak continuity*, Boll. Unione Mat. Ital. **9** (1974), 647–654.
- [4] T. NOIRI, A. AL-OMARI AND M.S.M. NOORANI, *On  $\omega b$ -open sets and  $b$ -Lindelöf spaces*, Eur. J. Pure Appl. Math. **1** (2008), 3–9.
- [5] S. WILLARD, *General topology*, Addison Wesley, London (1970).
- [6] S. WILLARD AND U.N.B. DISSANAYAKE, *The almost Lindelöf degree*, Canadian Math. Bull. **27** (1984), 452–455.
- [7] U. SENGUL, *Weakly  $b$ -continuous functions*, Chaos Solitons Fractals **41** (2009), 1070–1077.

Author's address:

Jamal M. Mustafa  
 Faculty of Science, Department of Sciences  
 Jerash University  
 Jerash, Jordan  
 E-mail: jjmrr971@yahoo.com

Received February 6, 2011  
 Revised March 21, 2011